arXiv:0907.4448v2 [math.AP] 6 Dec 2009

THE FRACTIONAL HARDY INEQUALITY WITH A REMAINDER TERM

BARTŁOMIEJ DYDA

Abstract. We calculate the regional fractional Laplacian on some power function on an interval. As an application, we prove Hardy inequality with an extra term for the fractional Laplacian on the interval with the optimal constant. As a result, we obtain the fractional Hardy inequality with best constant and an extra lower-order term for general domains, following the method developed by M. Loss and C. Sloane [11].

1. Main result and discussion

The purpose of this note is to prove the Hardy-type inequality with an extra term in the interval $(-1,1)$ – Corollary 2 – by the method of [5]. We will obtain it from the following formula analogous to the 'ground state representation' [8].

Theorem 1. *Let $0<\alpha<2$. Let $w(x)=(1-x^2)^{(\alpha-1)/2}$. For every $u\in C_c(-1,1)$,*

$$\begin{aligned}\mathcal{E}(u) &:= \frac{1}{2}\int_{-1}^{1}\int_{-1}^{1}\frac{(u(x)-u(y))^2}{|x-y|^{1+\alpha}}\,dx\,dy\\ &= \frac{1}{2}\int_{-1}^{1}\int_{-1}^{1}\left(\frac{u(x)}{w(x)}-\frac{u(y)}{w(y)}\right)^2\frac{w(x)w(y)}{|x-y|^{1+\alpha}}\,dx\,dy\\ &\quad + 2^\alpha\kappa_{1,\alpha}\int_{-1}^{1}u^2(x)\,(1-x^2)^{-\alpha}dx+\frac{1}{\alpha}\int_{-1}^{1}u^2(x)\,\phi(x)(1-x^2)^{-\alpha}\,dx; \qquad (1)\end{aligned}$$

where $\phi(x)=2^\alpha-(1+x)^\alpha-(1-x)^\alpha$ and

$$\kappa_{n,\alpha}=\pi^{\frac{n-1}{2}}\frac{\Gamma(\frac{1+\alpha}{2})}{\Gamma(\frac{n+\alpha}{2})}\frac{B\left(\frac{1+\alpha}{2},\frac{2-\alpha}{2}\right)-2^\alpha}{\alpha 2^\alpha}\,. \qquad (2)$$

Here B is the Euler beta function, and $C_c(a,b)$ denotes the class of all the continuous functions $u\colon(a,b)\to\mathbb{R}$ with compact support in (a,b).

Corollary 2. *Let $1<\alpha<2$. For every $u\in C_c(a,b)$,*

$$\begin{aligned}\frac{1}{2}\int_a^b\int_a^b\frac{(u(x)-u(y))^2}{|x-y|^{1+\alpha}}\,dx\,dy &\ge \kappa_{1,\alpha}\int_a^b u^2(x)\,\left(\frac{1}{x-a}+\frac{1}{b-x}\right)^\alpha dx\\ &\quad+\frac{4-2^{3-\alpha}}{\alpha(b-a)}\int_a^b u^2(x)\,\left(\frac{1}{x-a}+\frac{1}{b-x}\right)^{\alpha-1}dx. \qquad (3)\end{aligned}$$

Constant $\kappa_{1,\alpha}$ in (3) may not be replaced by a bigger constant.

2010 Mathematics Subject Classification: Primary 26D10, secondary 46E35, 31C25.
Key words and phrases: fractional Hardy inequality, best constant, interval, fractional Laplacian, censored stable process, convex domain, error term, ground state representation.
Research partially supported by grant MNiSW 3971/B/H03/2009/37.

On the right-hand side of (3) we note the function $\left(\frac{1}{x-a}+\frac{1}{b-x}\right) \geq \operatorname{dist}(x,(a,b)^c)$, where $\operatorname{dist}(x,(a,b)^c)$ equals the distance of x to the complement of (a,b). Analogous Hardy inequalities, involving the distance to the complement of a domain, and with optimal constants were derived in [7, 8, 4, 6, 11].

In fact, by the method developed in the paper [11], from Corollary 2 one gets the fractional Hardy inequality with a remainder term for general domains.

Theorem 3. *Let $1 < \alpha < 2$ and let $\Omega \subset \mathbb{R}^n$ be a bounded domain. For any $u \in C_c(\Omega)$*

$$\begin{aligned}\frac{1}{2}\int_{\Omega\times\Omega} \frac{(u(x)-u(y))^2}{|x-y|^{n+\alpha}}\,dx\,dy \geq{}& \kappa_{n,\alpha}\int_\Omega \frac{u^2(x)}{M_\alpha(x)^\alpha}\,dx\\ &+\frac{\pi^{(n-1)/2}\Gamma(\frac{\alpha}{2})(4-2^{3-\alpha})}{\alpha\Gamma(\frac{n+\alpha-1}{2})}\,\frac{1}{\operatorname{diam}\Omega}\int_\Omega \frac{u^2(x)}{M_{\alpha-1}(x)^{\alpha-1}}\,dx,\end{aligned}\tag{4}$$

where M_β is defined in [11, formula (6)]. *In particular, if Ω is a bounded convex domain, then for any $f \in C_c(\Omega)$ we have*

$$\begin{aligned}\frac{1}{2}\int_{\Omega\times\Omega} \frac{(u(x)-u(y))^2}{|x-y|^{n+\alpha}}\,dx\,dy \geq{}& \kappa_{n,\alpha}\int_\Omega \frac{u^2(x)}{\operatorname{dist}(x,\Omega^c)^\alpha}\,dx\\ &+\frac{\pi^{(n-1)/2}\Gamma(\frac{\alpha}{2})(4-2^{3-\alpha})}{\alpha\Gamma(\frac{n+\alpha-1}{2})}\,\frac{1}{\operatorname{diam}\Omega}\int_\Omega \frac{u^2(x)}{\operatorname{dist}(x,\Omega^c)^{\alpha-1}}\,dx,\end{aligned}\tag{5}$$

Constant $\kappa_{n,\alpha}$ in (4) *and* (5) *may not be replaced by a bigger constant.*

Finally, let us note that the symmetric bilinear form obtained from $\mathcal{E}$ by polarisation is, up to some multiplicative constant, the Dirichlet form of the censored stable process in $(-1,1)$, for the definition of its domain see [1]. We may also write a killed process counterparts of Theorem 1 and Corollary 2, and it is interesting that they have then a simpler form. Namely, we get the following

Corollary 4. *Let $0 < \alpha < 2$ and $w(x) = (1-x^2)^{(\alpha-1)/2}$. For every $u \in C_c(-1,1)$,*

$$\begin{aligned}\frac{1}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} \frac{(u(x)-u(y))^2}{|x-y|^{1+\alpha}}\,dx\,dy ={}&\frac{1}{2}\int_{-1}^1\int_{-1}^1 \left(\frac{u(x)}{w(x)}-\frac{u(y)}{w(y)}\right)^2 \frac{w(x)w(y)}{|x-y|^{1+\alpha}}\,dx\,dy\\ &+\frac{B(\frac{1+\alpha}{2},\frac{2-\alpha}{2})}{\alpha}\int_{-1}^1 u^2(x)\,(1-x^2)^{-\alpha}dx\\ \geq{}&\frac{B(\frac{1+\alpha}{2},\frac{2-\alpha}{2})}{\alpha 2^\alpha}\int_{-1}^1 u^2(x)\,\left(\frac{1}{x+1}+\frac{1}{1-x}\right)^\alpha dx.\end{aligned}$$

The constant $\frac{B(\frac{1+\alpha}{2},\frac{2-\alpha}{2})}{\alpha 2^\alpha}$ in the above inequality may not be replaced by a bigger one, and there is no remainder term of the form (3).

2. PROOFS

We define

$$Lu(x) = \lim_{\varepsilon\to 0^+}\int_{(-1,1)\cap\{|y-x|>\varepsilon\}} \frac{u(y)-u(x)}{|x-y|^{d+\alpha}}\,dy.$$

Note that L is, up to the multiplicative constant, the regional fractional Laplacian for an interval $(-1,1)$, see [9].

Lemma 5. *Let* $p > -1$ *and* $u_p(x) = (1-x^2)^p$. *For* $0 < \alpha < 2$ *we have*

$$Lu_p(x) = \frac{(1-x^2)^{p-\alpha}}{\alpha}\Bigg((1-x)^\alpha + (1+x)^\alpha - (2p+2-\alpha)B(p+1, 1-\alpha/2) + p.v.\int_{-1}^{1} \frac{(1-wx)^{\alpha-1-2p}-1}{|w|^{1+\alpha}}\,(1-w^2)^p\,dw\Bigg).$$

If we denote the principal value integral above by $I(p)$, *then*

$$I(\frac{\alpha-3}{2}) = x^2 B(p+1, 1-\alpha/2) \quad \text{if } 1<\alpha<2;$$
$$I(\frac{\alpha-2}{2}) = I(\frac{\alpha-1}{2}) = 0.$$

Proof. We have, by changing variable $w = y^2$ and then integrating by parts,

$$\begin{aligned}Lu_p(0) &= 2\lim_{\varepsilon\to0^+}\int_\varepsilon^1 \frac{(1-y^2)^p-1}{y^{1+\alpha}}\,dy \\ &= 2\lim_{\varepsilon\to0^+}\left(\frac12\int_{\varepsilon^2}^1 (1-w)^p w^{-1-\alpha/2}[(1-w)+w]\,dw - \int_\varepsilon^1 y^{-1-\alpha}\,dy\right) \\ &= 2\lim_{\varepsilon\to0^+}\Bigg(\frac1\alpha(1-\varepsilon^2)^{p+1}\varepsilon^{-\alpha} - \frac{p+1}{\alpha}\int_{\varepsilon^2}^1(1-w)^p w^{-\alpha/2}\,dw \\ &\qquad + \frac12\int_{\varepsilon^2}^1 (1-w)^p w^{-\alpha/2}\,dw + \frac1\alpha - \frac{\varepsilon^{-\alpha}}{\alpha}\Bigg)\end{aligned}$$

It is easy to see that

$$\lim_{\varepsilon\to0^+}\left(\frac1\alpha(1-\varepsilon^2)^{p+1}\varepsilon^{-\alpha} - \frac{\varepsilon^{-\alpha}}{\alpha}\right) = \lim_{\varepsilon\to0^+}\frac{\varepsilon^{2-\alpha}}{\alpha}\,\frac{(1-\varepsilon^2)^{p+1}-1}{\varepsilon^2} = 0.$$

Hence

$$Lu_p(0) = \frac{2}{\alpha}(1-(p+1-\alpha/2)B(p+1,1-\alpha/2)).$$

We have, for $x_0 \in (-1,1)$,

$$Lu_p(x_0) = p.v.\int_{-1}^1 \frac{(1-y^2)^p - (1-x_0^2)^p}{|y-x_0|^{1+\alpha}}\,dy.$$

We change the variable in the following way

$$\begin{aligned} w = \varphi(y) &:= \frac{x_0-y}{1-x_0y}; \qquad y = \varphi(w); \\ \varphi'(y) &= \frac{x_0^2-1}{(1-x_0y)^2}; \\ y - x_0 &= \frac{w(1-x_0^2)}{wx_0-1}; \\ 1-y^2 &= \frac{(1-x_0^2)(1-w^2)}{(wx_0-1)^2}. \end{aligned}$$

Moreover the principal value integral transforms well. We obtain

$$
\begin{aligned}
Lu_p(x_0) &= (1-x_0^2)^{p-\alpha} p.v. \int_{-1}^{1} \frac{(1-w^2)^p - (1-wx_0)^{2p}}{|w|^{1+\alpha}} (1-wx_0)^{\alpha-1-2p}\, dw \\
&= (1-x_0^2)^{p-\alpha}\Bigg[Lu_p(0) - p.v. \int_{-1}^{1} \frac{(1-wx_0)^{\alpha-1}-1}{|w|^{1+\alpha}}\, dw \qquad (6) \\
&\quad + p.v. \int_{-1}^{1} \frac{(1-wx_0)^{\alpha-1-2p}-1}{|w|^{1+\alpha}} (1-w^2)^p\, dw \Bigg].
\end{aligned}
$$

We calculate the integral in (6)

$$
I := p.v. \int_{-1}^{1} \frac{(1-wx_0)^{\alpha-1}-1}{|w|^{1+\alpha}}\, dw = \lim_{\varepsilon\to 0^+} (J_\varepsilon(x_0) + J_\varepsilon(-x_0)),
$$

where

$$
\begin{aligned}
J_\varepsilon(x_0) &= \int_{\varepsilon}^{1} \frac{(1-wx_0)^{\alpha-1}-1}{w^{1+\alpha}}\, dw \\
&= \int_{\varepsilon}^{1} \Big(\frac{1}{w} - x_0\Big)^{\alpha-1} \frac{dw}{w^2} - \frac{\varepsilon^{-\alpha}-1}{\alpha} \\
&= \frac{1}{\alpha}\Big(\frac{1}{\varepsilon} - x_0\Big)^{\alpha} - \frac{1}{\alpha}\Big(1-x_0\Big)^{\alpha} - \frac{\varepsilon^{-\alpha}-1}{\alpha} \\
&= \frac{1}{\alpha} - \frac{1}{\alpha}\Big(1-x_0\Big)^{\alpha} + \frac{(1-\varepsilon x_0)^{\alpha}-1}{\alpha\varepsilon^{\alpha}}.
\end{aligned}
$$

By l'Hôspital rule we find that

$$
I = \frac{2}{\alpha} - \frac{1}{\alpha}\Big(1-x_0\Big)^{\alpha} - \frac{1}{\alpha}\Big(1+x_0\Big)^{\alpha},
$$

and the first part of the lemma is proved.

Obtaining $I(p)$ for the three values of p is easy and is omitted. □

Next lemma is in fact a very special case of Proposition 2.3 from [6]. In this special case the proof may be simplified, and for reader's convenience we give the sketch of the proof.

Lemma 6. *For every* $u \in C_c(-1,1)$ *and any strictly positive function* $w \in C^2(-1,1)$ *such that* $Lw \le 0$, *we have*

$$
\mathcal{E}(u) = \int_{-1}^{1} u^2(x) \frac{-Lw(x)}{w(x)}\, dx + \frac{1}{2}\int_{-1}^{1}\int_{-1}^{1} \left(\frac{u(x)}{w(x)} - \frac{u(y)}{w(y)}\right)^2 \frac{w(x)w(y)}{|x-y|^{1+\alpha}}\, dx\, dy.
$$

Proof. We use the elementary equality [4, (9)]

$$
\begin{aligned}
(u(x)-u(y))^2 + u^2(x)\frac{w(y)-w(x)}{w(x)} + u^2(y)\frac{w(x)-w(y)}{w(y)} \\
= \left(\frac{u(x)}{w(x)} - \frac{u(y)}{w(y)}\right)^2 w(x)w(y),
\end{aligned}
$$

then integrate against the measure $1_{\{|x-y|>\varepsilon\}}|x-y|^{-1-\alpha}\, dx\, dy$ and take $\varepsilon \to 0$. We use Taylor expansion for w and the compactness of the support of u to apply Lebesgue dominated convergence theorem. □

Proof of Theorem 1. The theorem follows immediately from Lemma 5 applied to $w(x) = (x^2-1)^{(\alpha-1)/2}$ and Lemma 6 applied to $p = (\alpha-1)/2$. $\square$

Proof of Corollary 2. By Theorem 1 and symmetry of the function ϕ, to prove (3) it is enough to show that

$$\phi(x) := 2^\alpha - (1+x)^\alpha - (1-x)^\alpha \geq (2^\alpha - 2)(1-x^2), \quad x \in [0,1]. \tag{7}$$

After substitution $u = x^2$, we see that it is enough to show that the function

$$g(u) := (2^\alpha - 2)u - (1-\sqrt{u})^\alpha - (1+\sqrt{u})^\alpha + 2$$

is concave. To show this, it suffices to show that

$$g'(u) = 2^\alpha - 2 + \frac{\alpha}{2\sqrt{u}}\left((1-\sqrt{u})^{\alpha-1} - (1+\sqrt{u})^{\alpha-1}\right)$$

is a decreasing function. We again substitute $t = u^2$ and observe that

$$\frac{(1-t)^{\alpha-1} - (1+t)^{\alpha-1}}{t} = \frac{h(t) - h(0)}{t},$$

where $h(t) = (1-t)^{\alpha-1} - (1+t)^{\alpha-1}$. Now since h is concave, the function $t \mapsto \frac{h(t)-h(0)}{t}$ is decreasing, and so is the function g'. This proves the claim (7) and, consequently, the inequality (3).

The fact that the constant $\kappa_{1,\alpha}$ in (3) is optimal follows from [4]. $\square$

Proof of Theorem 3. The proof is analogous to the proof of Theorem 1.1 in [11]. In inequality [11, (33)] we would get an extra lower order term. Namely, using notation from [11], the extra term would be

$$\begin{aligned}
&\frac{1}{2}\lambda_{1,\alpha}\int_{S^{n-1}} dw \int_{\{x:x\cdot w=0\}} d\mathcal{L}_w(x) \int_{x+sw\in\Omega} ds|f(x+sw)|^2 \\
&\qquad\qquad \times \left[\frac{1}{d_w(x+sw)} + \frac{1}{\delta_w(x+sw)}\right]^{\alpha-1} \frac{1}{d_w(x+sw)+\delta_w(x+sw)} \\
&= \frac{1}{2}\lambda_{1,\alpha}\int_{S^{n-1}} dw \int_\Omega |f(x)|^2 \left[\frac{1}{d_{w,\Omega}(x)} + \frac{1}{\delta_{w,\Omega}(x)}\right]^{\alpha-1} \frac{1}{d_{w,\Omega}(x)+\delta_{w,\Omega}(x)} \\
&\qquad\qquad \geq \frac{\lambda_{n,\alpha}}{\operatorname{diam}\Omega}\int_\Omega \frac{|f(x)|^2}{M_{\alpha-1}(x)^{\alpha-1}}\,dx.
\end{aligned}$$

Thus we get (4).

Note that [11, (9)] is valid for any $\alpha > 0$. We apply it to $\alpha - 1$ in place of α and we get inequality (5). $\square$

Proof of Corollary 4. The equality follows from Theorem 1 and the following easy formula

$$\frac{1}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} \frac{(u(x)-u(y))^2}{|x-y|^{1+\alpha}}\,dx\,dy = \frac{1}{2}\mathcal{E}(u) + \int_{-1}^{1} u^2(x)\frac{1}{\alpha}\left((1+x)^{-\alpha} + (1-x)^{-\alpha}\right)dx.$$

The sharpness of the constant and the lack of a remainder term follow from approximating the function w, for example by $u = \psi_n w$, where $\operatorname{supp}\psi_n \subset [-1+1/n, 1-1/n]$, $\psi_n(x) = 1$ on $(-1+2/n, 1-2/n)$ and $|\psi_n'(x)| \leq 2n$ on $(-1,1)$. $\square$

REFERENCES

[1] K. Bogdan, K. Burdzy, and Z.-Q. Chen. Censored stable processes. *Probab. Theory Related Fields*, 127(1):89–152, 2003.

[2] Z.-Q. Chen and R. Song. Hardy inequality for censored stable processes. *Tohoku Math. J. (2)*, 55(3):439–450, 2003.

[3] B. Dyda. A fractional order Hardy inequality. *Ill. J. Math.*, 48(2):575–588, 2004.

[4] B. Dyda and K. Bogdan. The best constant in a fractional hardy inequality. *Math. Nachr.*

[5] P. J. Fitzsimmons. Hardy's inequality for Dirichlet forms. *J. Math. Anal. Appl.*, 250(2):548–560, 2000.

[6] R. Frank and R. Seiringer. Sharp fractional Hardy inequalities in half-spaces. arXiv:0906.1561v1 [math.FA], 2009.

[7] R. L. Frank, E. H. Lieb, and R. Seiringer. Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators. *J. Amer. Math. Soc.*, 21(4):925–950, 2008.

[8] R. L. Frank and R. Seiringer. Non-linear ground state representations and sharp Hardy inequalities. *J. Funct. Anal.*, 255(12):3407–3430, 2008.

[9] Q.-Y. Guan and Z.-M. Ma. Reflected symmetric α-stable processes and regional fractional Laplacian. *Probab. Theory Related Fields*, 134(4):649–694, 2006.

[10] N. Krugljak, L. Maligranda, and L. E. Persson. On an elementary approach to the fractional Hardy inequality. *Proc. Amer. Math. Soc.*, 128(3):727–734, 2000.

[11] M. Loss and C. Sloane. Hardy inequalities for fractional integrals on general domains. arXiv:0907.3054v1 [math.AP], 2009.

INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE, WROCŁAW UNIVERSITY OF TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCŁAW, POLAND

E-mail address: `bdyda@pwr.wroc.pl`